

Should we exclude children with overweight or obesity from height reference samples?


Mathieu Roelants¹, Bente Brannsether², Hege Kristiansen³,
Robert Bjercknes⁴, Karel Hoppenbrouwers¹, Pétur B Júlíusson^{4,5}

¹ Department of Public Health and Primary Care, KULeuven - University of Leuven, Belgium; ² Department of Pediatrics, Stavanger University Hospital, Norway; ³ Department of Pediatrics, District General Hospital of Førde, Norway; ⁴ Department of Clinical Science, University of Bergen, Norway; ⁵ Department of Pediatrics, Haukeland University Hospital, Bergen, Norway



Introduction

- Growth references show how children grow in a particular population (genetic background), at a particular time (changing environment)
- Used to monitor overall health and wellbeing
- Children with conditions known to affect growth usually excluded
- Overweight/obese children:
 - Height also different
 - Increasing prevalence →
 - **Also exclude?**



Materials and Methods (1)

- Cross-sectional data from 2 growth reference studies
- 12252 Belgian and 6159 Norwegian children
- 2 – 18 years of age
- Measured (height, weight → BMI) in 2002–2006
- Weight status defined by IOTF¹:

not overweight, overweight (excl. obese), obese

¹Cole et al. BMJ, 2001

Materials and Methods (2)

1) Is height different in these “*weight classes*” ?

- height (cm) → SD scores
- anova within age groups (both sexes combined)

2) Are height-by-age references affected?

- Three height reference curves estimated:
 - All children (= *current national reference*)
 - Excluding *obese* children only
 - Excluding *overweight* and *obese* children
- LMS method², same # edf

²Cole and Green, Stat. Med. 1992

KU LEUVEN

Results (1)


- Height SDS according to weight status³

AGE	Not overweight			Overweight (excl. obese)			Obese			anova
	n	mean	SDS (95%CI)	n	mean	SDS (95%CI)	n	mean	SDS (95%CI)	
Belgium										
2 – 5	2594	-0.04	(-0.08 – 0.00)	213	0.36	(0.21 – 0.51)	47	0.30	(0.00 – 0.61)	p < 0.001
6 – 8	2122	-0.08	(-0.12 – -0.03)	227	0.49	(0.35 – 0.62)	81	0.65	(0.42 – 0.88)	p < 0.001
9 – 11	2331	-0.09	(-0.13 – -0.05)	338	0.38	(0.27 – 0.48)	80	0.73	(0.52 – 0.94)	p < 0.001
12 – 14	2010	-0.04	(-0.08 – 0.01)	258	0.21	(0.09 – 0.33)	70	0.36	(0.12 – 0.60)	p < 0.001
15 – 17	1574	0.02	(-0.03 – 0.07)	199	0.02	(-0.12 – 0.15)	56	-0.03	(-0.35 – 0.30)	p = 0.3
Norway										
2 – 5	1151	-0.05	(-0.11 – 0.01)	148	0.14	(-0.02 – 0.30)	18	0.49	(-0.08 – 1.07)	p < 0.001
6 – 8	989	-0.12	(-0.18 – -0.06)	149	0.43	(0.27 – 0.59)	56	0.72	(0.44 – 0.99)	p < 0.001
9 – 11	944	-0.09	(-0.16 – -0.03)	162	0.42	(0.27 – 0.56)	29	0.50	(0.08 – 0.91)	p < 0.001
12 – 14	1074	-0.05	(-0.11 – 0.02)	121	0.12	(-0.09 – 0.32)	13	0.24	(-0.23 – 0.70)	p < 0.001
15 – 17	1130	-0.01	(-0.07 – 0.05)	129	-0.09	(-0.26 – 0.08)	26	-0.15	(-0.57 – 0.27)	p = 0.6

³prevalence of overweight (incl. obesity) and obesity = 13.0% and 2.8% in the Belgian sample and 13.8% and 2.3% in the Norwegian sample.

Results (2)

- Effect of exclusion on the growth chart




(standard curve) - (new curve)

All children = std: $\Delta=0$

Excluding ob < std

Excluding ow + ob << std



Results (3)

- Theoretical referral rates (all children, different reference):

Reference based on:	Percentage (95%CI) below -2SD			Percentage (95%CI) above + 2SD		
	All	Non overweight/ obese	Non obese	All	Non overweight/ obese	Non obese
Belgium						
2-5	2.2 (1.7 – 2.9)	2.2 (1.7 – 2.8)	2.2 (1.7 – 2.9)	2.5 (2.0 – 3.2)	3.0 (2.4 – 3.7)	2.6 (2.0 – 3.3)
6-8	2.1 (1.6 – 2.8)	1.8 (1.3 – 2.4)	2.0 (1.5 – 2.7)	2.0 (1.5 – 2.6)	2.5 (1.9 – 3.2)	2.0 (1.5 – 2.7)
9-11	2.0 (1.5 – 2.6)	1.7 (1.3 – 2.3)	2.0 (1.5 – 2.6)	2.7 (2.1 – 3.4)	3.1 (2.5 – 3.8)	2.7 (2.2 – 3.4)
12-14	2.3 (1.7 – 3.0)	2.2 (1.6 – 2.9)	2.3 (1.7 – 3.0)	2.2 (1.6 – 2.9)	2.2 (1.7 – 2.9)	2.1 (1.6 – 2.8)
15-17	2.5 (1.8 – 3.3)	2.5 (1.9 – 3.4)	2.5 (1.9 – 3.4)	1.8 (1.3 – 2.6)	1.7 (1.2 – 2.4)	1.9 (1.3 – 2.6)
Norway						
2-5	2.7 (1.9 – 3.8)	2.6 (1.8 – 3.6)	2.7 (1.9 – 3.8)	2.4 (1.6 – 3.4)	2.8 (2.0 – 3.9)	2.5 (1.8 – 3.5)
6-8	2.3 (1.5 – 3.3)	2.0 (1.3 – 3.0)	2.2 (1.5 – 3.2)	2.1 (1.4 – 3.1)	3.3 (2.4 – 4.5)	2.6 (1.8 – 3.7)
9-11	1.5 (0.9 – 2.4)	1.3 (0.8 – 2.2)	1.5 (0.9 – 2.4)	2.5 (1.7 – 3.7)	3.3 (2.4 – 4.6)	2.8 (2.0 – 4.0)
12-14	3.0 (2.1 – 4.1)	2.9 (2.1 – 4.1)	3.0 (2.1 – 4.1)	2.2 (1.5 – 3.3)	2.6 (1.8 – 3.7)	2.3 (1.6 – 3.4)
15-17	2.1 (1.4 – 3.1)	2.1 (1.4 – 3.1)	2.1 (1.4 – 3.1)	2.3 (1.6 – 3.4)	2.3 (1.6 – 3.4)	2.3 (1.6 – 3.4)

Conclusion

- Typical height pattern confirmed:
 - Overweight and obese children are on average taller before puberty
 - Earlier maturity?
- Similar effect on the growth chart when excluding children with overweight or obesity
 - Effect of additional exclusion depends on *effect size* and *prevalence*: larger for overweight!
 - Could be larger in populations with higher prevalence
- Minor consequences for clinical cut-offs

Should we exclude children with overweight or obesity from height reference samples?

Mathieu Roelants¹, Bente Brannsether², Hege Kristiansen³,
Robert Bjerknes⁴, Karel Hoppenbrouwers¹, Pétur B Júlíusson^{4,5}

¹ Department of Public Health and Primary Care, KULeuven - University of Leuven, Belgium;

² Department of Pediatrics, Stavanger University Hospital, Norway; ³ Department of Pediatrics, District General Hospital of Førde, Norway; ⁴ Department of Clinical Science, University of Bergen, Norway; ⁵ Department of Pediatrics, Haukeland University Hospital, Bergen, Norway

Should we exclude children with overweight or obesity from height reference samples? Data from Belgium and Norway shows that it makes little difference!

Mathieu Roelants¹, Bente Brannsether², Hege Kristiansen³,
Robert Bjercknes⁴, Karel Hoppenbrouwers¹, Pétur B Júlíusson^{4,5}

¹ Department of Public Health and Primary Care, KULeuven - University of Leuven, Belgium;

² Department of Pediatrics, Stavanger University Hospital, Norway; ³ Department of Pediatrics, District General Hospital of Førde, Norway; ⁴ Department of Clinical Science, University of Bergen, Norway;

⁵ Department of Pediatrics, Haukeland University Hospital, Bergen, Norway